

**MACROZOOBENTHOS OF LAKES OF UZBEKISTAN****Matmurotov Muratbay Allamurotovich**  
teacherKarakalpak State University,  
Nukus, Uzbekistan**Saparov Abdurahman Danabaevich**candidate of biological sciences, dean  
Nukus State Pedagogical Institute,  
Nukus, Uzbekistan**Ismoilov Khojiakbar Faizievich**assistant  
Tashkent State Agrarian University,  
Tashkent, Uzbekistan**Beisheeva Shahnoza Azadovna**teacher  
Secondary school No. 71,  
Tashkent, Uzbekistan**Mirzambetov Nurlibay**trainee  
Karakalpak Research Institute of Natural Sciences,  
Nukus, Uzbekistan**ДОННАЯ ФАУНА НЕКОТОРЫХ ОЗЕР УЗБЕКИСТАНА****Матмуратов Муратбай Алламуратович**  
преподавательКаракалпакский государственный университет,  
г. Нукус, Узбекистан**Сапаров Абдурахман Данабаевич**кандидат биологических наук, декан факультета  
Нукусский государственный педагогический университет,  
г. Нукус, Узбекистан**Исмоилов Ходжаакбар Фаизиевич**ассистент  
Ташкентский государственный аграрный университет,  
г. Ташкент, Узбекистан**Бейшеева Шахноза Азадовна**преподаватель  
Общеобразовательная средняя школа № 71,  
г. Ташкент, Узбекистан**Мирзамбетов Нурлыбай**стажер  
Каракалпакский научно-исследовательский институт естественных наук,  
Нукус, УзбекистанDOI: [10.31618/nas.2413-5291.2020.3.58.279](https://doi.org/10.31618/nas.2413-5291.2020.3.58.279)**Abstract**

Species composition of macrozoobenthos of 6 lakes from different regions (north, center, south) of the republic have been studied. Total 111 species of bottom animals have been recorded. List of dominant species is presented. Using coefficient of Sørensen-Czekanowski similarity of species composition of macrozoobenthos of the lakes studied have been determined.

**Резюме**

Изучен качественный состав макрозообентоса 6 озер из разных регионов (север, центр, юг) республики. Всего отмечен 111 вид донных животных. Приведен список доминирующих видов. При помощи коэффициента Сёренсена-Чекановского определено сходство качественного состава макрозообентоса исследованных озер.

**Keywords:** macrozoobenthos, lakes, species composition, fauna similarity, Uzbekistan.

**Ключевые слова:** макрозообентос, озера, видовой состав, сходство фаун, Узбекистан.

**Introduction**

The biological productivity of water bodies is determined by the vital activity of hydrobionts, one of the most important link of which in aquatic ecosystem

of water bodies is the aquatic bottom fauna (zoobenthos). Benthic animals enter into various, primarily trophic, connections with organisms of practically all ecotopes of water bodies. The most

important role of macrozoobenthos organisms play as elements of the fish food. Such important commercial fish species as the carp, crucian carp, bream, roach, black amur, and also such Red Data Book species as barbell, shovel-noses, white-eye bream feed largely on aquatic benthic fauna [5]. In the waterbodies of Uzbekistan and throughout Central Asia, the bottom fauna makes up the bulk of the forage base of most commercial fish species.

In Uzbekistan the macrozoobenthos of lakes was studied by A.M. Mukhamediev [6], S. Embergenov [2, etc.], B. Bekmurzaev [1, etc.] and other. However, most of these works date back to the 1960-1970s. Also, there are practically no generalizing and comparative works.

#### Material and methods

The material is collected seasonally (spring, summer, autumn) during 2000-2018 using Petersen's dredging and is processed according to conventional methods [9-12]. Identification of hydrobionts led by modern guides [7, 8, 13, etc.].

To assess the similarities and differences in the species composition of the macrozoobenthos of lakes, the Sørensen-Czekanowski formula was used:

$$K=2C \times 100\%$$

$$(A+B)$$

where A and B are number of species in two compared lakes, C is the number of species common to this pair, K is the coefficient of similarity of the species composition.

Seven lakes from different (south, north, west, east, center) regions of the republic were studied: Sarykamysh (Karakalpakstan), Western Karateren (Wetland Sudochie, Karakalpakstan), Ullishorkul (Khorezm viloyat = province), Karakir (Bukhara viloyat), Tuzkan (Jizzakh viloyat), Sarykamysh (Fergana viloyat) (fig. 1).



Fig. 1. Location of the lakes studied. 1: Sarykamysh, 2: Western Karateren, 3: Ullishorkul, 4: Karakir, 5: Tuzkan, 6: Sarykamysh, 7: Sichankul

#### Results and discussion

As can be seen from the table 1, the studied lakes of different areas, depths, water salinity, with different

soils. The only freshwater lake is Sarykamysh in the Fergana Valley. The rest of the lakes are brackish.

Table 1.

Main abiotic parameters of the studied lakes

Lakes	Area, 10 <sup>3</sup> ha	Maximal depths, m	Dominant depths, m	Salinity, g/l	Soils
Sarykamysh (Karakalpakstan)	387.5	40.0	5.0-7.0	11.0-13.0	silt and sand
West Karateren	0.4	1.7	0.8-1.2	4.0-6.0	gray silt
Tuzkan	700.0	20.0	12.0	10.0	gray silt
Sarykamysh (Fergana viloyat)	26.2	7.0	3.0	0.8	gray and black silts

Ullishorkul	1.8	4.5	1.5	5.8	gray silt
Karakir	26.2	5.0	1.5-2.0	10.7	sand
Sichankul	7.5	20.0	9.0	7.0	sand

A total of 187 species of bottom animals were observed, the most diverse was the benthos of freshwater lake Sarykamys (Fergana viloyat), the least - in brackish lakes (tables 2, 3). As in most of lakes in Central Asia, chironomids were most diverse (86 species = almost half of all species of

macrozoobenthos) (table 3). Other groups are represented by a much smaller number of species: Odonata – 15 species, Coleoptera – 7, Diptera (excluding chironomids) – 10, Mollusca – 12, Annelida – 10, Ephemeroptera – 11. The rest of the groups of organisms are represented by single species.

Table 2.

**Dominant species of macrozoobenthos of studied lakes (SKK – Sarykamys, Karakalpakstan), KT – West Karateren, TZ – Tuzkan, SKF – Sarykamys (Fergana Valley), US – Ullishorkul, KK – Karakir, SI – Sichankul)**

Taxa / Lakes	SKK	KT	TZ	SKF	US	KK	SI
ANNELIDA							
<i>Paranis simplex</i> Hrabce	+	+	+	-	-	+	+
<i>P. littoralis</i> O.F. Müller	+	-	+	-	-	+	+
<i>Tubifex sp.</i>	-	+	-	+	+	-	-
<i>Nereis diversicolor</i> O.F. Müller	+	+	-	-	-	-	-
MOLLUSCA							
<i>Colletopterum cyreum</i> (Kobelt)	-	-	-	+	-	-	-
<i>Corbicula fluminalis</i> O.F. Müller	-	-	-	+	-	-	-
<i>Physella acuta</i> Draparnaud	-	+	+	+	-	-	+
<i>Cerastoderma isthmicum</i> Issel	+	-	-	-	-	-	-
<i>Lymnaea truncatula</i> O.F. Müller	-	+	+	+	-	-	-
<i>Caspihydrobia conica</i> Logvin. et Starobog.	-	+	-	-	-	-	-
CRUSTACEA							
<i>Macrobrachium nipponense</i> De Haan	+	+	+	+	+	+	+
<i>Turkogammarus aralensis</i> (Uljanin)	+	+	-	-	-	-	-
<i>Mesomysis kowalevskii</i> Czerniavsky	-	-	+	+	+	-	+
<i>Paramysis lacustris</i> (Czerniavsky)	+	+	-	+	+	-	+
INSECTA							
<i>Anax imperator</i> Leach	-	-	-	-	-	+	+
<i>Cloen dipterum</i> L.	-	-	+	+	+	+	-
<i>Caenis macrura</i> Stephens	-	+	+	+	+	-	-
<i>Ecnomus tenellus</i> Rambur	-	-	+	+	-	-	-
<i>Chironomus salinarius</i> Kieffer	+	+	+	-	+	+	+
<i>Ch. Halophilus</i> Kieffer	+	+	-	-	-	+	+
<i>Ch. thummi</i> Kieffer	+	+	+	+	+	+	-
<i>Cricotopus tenellus</i> Fabricius	-	-	-	+	-	-	-
<i>C. silvestris</i> (Fabricius)	-	+	+	+	+	+	+
<i>Procladius ferrugineus</i> Kieffer	+	+	+	+	+	+	-
<i>Polypedilum aberrans</i> Tschern.	+	+	-	+	+	-	+
<i>Endochironomus tendens</i> (Fabricius)	+	-	-	+	+	-	-
<i>Glyptotendipes barbipes</i> (Staeger)	-	-	-	+	-	-	-
<i>G. glaucus</i> (Meigen)	-	-	-	+	-	-	-
Total number of species	33	30	25	75	28	28	31
Lakes	SKK	KT	TZ	SKF	US	KK	SI

The remains of the Aral Sea aquatic fauna have been preserved in the lakes of Sarykamys (Karakalpakstan) and Western Karateren: polychaete *Nereis diversicolor*, mollusks *Cerastoderma*

*isthmicum*, *Caspihydrobia cf. conica*, *Theodoxus pallasi*, amphipod *Turkogammarus aralensis* (included in the Red Data Book of the Republic of Uzbekistan [3]).

Table 3.

**Taxonomic diversity (number of species) of macrozoobenthos of the lakes studied (legend as in table 2)**

Taxa / Lakes	SKK	KT	TZ	SKF	US	KK	SI	Number of species / %
Annelida	5	4	3	6	4	4	5	10 / 5.3
Mollusca	3	1	1	7	1	1	1	12 / 6.4
Crustacea	2	1	1	1	1	1	3	5 / 2.7
Chironomidae	20	19	17	49	18	18	19	86 / 46.0
Other Diptera	5	5	4	5	9	5	5	15 / 8.0
Odonata	4	3	2	8	3	3	4	15 / 8.0
Coleoptera	1	2	2	3	1	1	2	7 / 3.7
Ephemeroptera	2	2	2	7	3	2	3	11 / 5.9
Trichoptera	1	2	1	4	2	1	2	6 / 3.2
Hemiptera	3	2	3	5	2	2	2	9 / 4.8
Plecoptera	2	2	2	4	3	3	2	7 / 3.7
Hydracarina	1	1	1	2	2	1	1	4 / 2.1
Total species	49	44	39	101	49	42	49	187

Comparison of the species composition of zoobenthos in the studied lakes revealed that the species composition of brackish water lakes is the most similar. The species composition of the

macrozoobenthos of the freshwater lake Sarykamys of the Fergana Valley is sharply different from the species composition of the brackish lakes studied (table 4) and more similar to the bottom fauna of reservoirs [4].

Table 4.

**The similarity of the species composition of macrozoobenthos of the lakes studied according to the Sørensen-Czekanowski coefficient (legend as in table 2)**

Lakes	SKK	KT	TZ	KK	US	SKF	SI
Sarykamys (Karakalpakstan)		80	75	69	77	20	77
Western Karateren	80		68	70	78	21	66
Tuzkan	75	68		75	71	18	70
Karakir	81	70	75		68	19	67
Ullishorkul	77	78	71	68		22	56
Sarykamys (Fergana)	20	21	18	19	22		23
Sichankul	77	66	70	67	56	23	

As practical recommendations, it is possible to propose the introduction of some invertebrates of the Aral Sea fauna such as *Nereis diversicolor*, *Turkogammarus aralensis*, *Theodoxus pallasi*, and *Cerastoderma isthmicum* into the brackish-water lakes Ullishorkul, Karakir, Tuzkan, Sichankul, and other brackish waterbodies of the republic.

**Acknowledgements**

The authors are grateful to O.D. Gerasimova for help in identifying of some species and to I.M. Mirabdullayev for valuable comments and editing the manuscript.

**References**

1. Bekmurzaev B. Benthos Karazharskoi sistemy ozer delty Amudari [Benthos of the Karazhar Lake System in Amudarya river delta] // Uzbek biol. Zhurn. 1969. 2 (in Russian).
2. Embergenov S., Khusainva N.Z. O zoobentose ozera Vostochnyi Karateren (delta Amudarii) [On the zoobenthos of lake Eastern Karateren (Amudarya river delta)] // Vestnik Karakalpak. Filiala AN UzSSR. 1970. № 2 (in Russian).
3. Krasnaya Kniga Respubliki Uzbekistan [The Red Data Book of the Republic of Uzbekistan]. – Tashkent: Chinor ENK, 2019. V. 2. (in Russian).

4. Matmuratov M.A., Abdinazarov H.Kh., Sabirov Zh.Zh. et al. Makrozoobentos nekotorykh ravninnykh vodokhranilishch Uzbekistana [Macrozoobenthos of some flat reservoirs of Uzbekistan] // Vestnik Karakalpak. Otdel. Akad. Nauk Resp. Uzb. 2016. № 3. (in Russian).

5. Mirabdullayev I.M., Mirzaev U.T., Kuzmetov A.R., Kimsanov Z.O. Uzbekistan va kushni hududlar baliklar aniklagichi [Guide to fish of Uzbekistan and adjacent territories]. – Tashkent: Sano-Standart, 2011. (in Uzbek).

6. Mukhamediev A.M. Gidrobiologiya vodoemov Ferganskoi doliny [Hydrobiology of waterbodies of the Fergana Valley]. – Tashkent: Fan, 1967. – (in Russian).

7. Opredelitel zooplanktona i zoobentosa presnykh vod Evropeiskoi Rossii [Key to zooplankton and zoobenthos freshwaters of European Russia]. V. 2. Zoobenthos. – M. - SPb.: Tovar. nauch. izdat. KMK, 2016. (in Russian).

8. Opredelitel presnovodnykh bespozvonochnykh Rossii i sopredelnykh territoriy [Key to freshwater invertebrates of Russia and adjacent lands]. – SPb.: Zool Inst. RAS, 1994-2004. V. 1-6. (in Russian).

9. Plotnikov G.K., Peskova T.E., Shkute A., Pupinya A., Pupinsh M. Sbornik klassicheskikh metodov gidrobiologicheskikh issledovaniy dlya ispolzovaniy v akvakulture [Collection of classic

methods in hydrobiological researches for using in aquaculture]. – Daugavpils: Saule, 2017. (in Russian).

10. Popchenko V.I., Bulgakov G.P. Monitoring makrozoobentosa [Monitoring of macrozoobenthos] // In: Rukovodstvo po gidrobiologicheskomu monitoringu presnovodnykh ekosistem. SPb.: Gidrometeoizdat, 1992. (in Russian).

11. Prakticheskaya gidrobiologiya. Presnovodnye ekosistemy [Practical hydrobiology. Freshwater ecosystems] (Eds. V.D. Fedorov, V.I. Kapkov). – M.: Izd. MSU, 2006. (in Russian).

12. Salazkin A.A., Alimov A.F., Finogenova N.P. Metodicheskie rekomendatsii po sboru i obrabotke

materialov pri gidrobiologicheskikh issledovaniyakh na presnovodnykh vodoyemakh: zoobentos i ego produktsiya [Methodical recommendations for collecting and processing of materials under hydrobiological researches on freshwater waterbodies: zoobenthos and its production]. – Leningrad: GosNIORKh, 1984. (in Russian).

13. Gloer R, Meier-Brook C. Süßwassermollusken. Ein Bestimmungsschlüssel für die Bundesrepublik Deutschland. – Hamburg: Deutsch. Jugendbund Naturbeobacht., 2003.